

- 1 a Show that the equation $x^3 - 7x - 11 = 0$ has a real root in the interval (3, 4).
 b Using the iterative formula $x_{n+1} = \sqrt{7 + \frac{11}{x_n}}$, with $x_0 = 3.2$, find x_1, x_2 and x_3 , giving the value of x_3 correct to 2 decimal places.


2
$$f(x) \equiv 4 \operatorname{cosec} x - 5 + 2x.$$

- a Find the values of $f(4)$ and $f(5)$.
 b Hence show that the equation $f(x) = 0$ has a root in the interval (4, 5).

The iterative formula $x_{n+1} = a + \frac{b}{\sin x_n}$, where a and b are constants, is used to find this root.

- c Find the values of a and b .
 d Starting with $x_0 = 4.5$, use the iterative formula with your values of a and b to find 3 further approximations of the root, giving your final answer correct to 3 decimal places.

3



The diagram shows the curve with equation $y = f(x)$ where

$$f : x \rightarrow 2x + \ln(3x - 1), \quad x \in \mathbb{R}, \quad x > \frac{1}{3}.$$

Given that $f(\alpha) = 0$,

- a show that $0.4 < \alpha < 0.5$,
 b use the iterative formula $x_{n+1} = \frac{1}{3}(1 + e^{-2x_n})$, with $x_0 = 0.45$, to find the value of α correct to 3 decimal places.
- 4 a On the same set of axes, sketch the curves $y = \cos x$ and $y = x^2$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$.
 b Show that the equation $\cos x - x^2 = 0$ has exactly one positive and one negative real root.
 c Show that the positive real root lies in the interval $[0.8, 0.9]$.
 d Use the iteration formula $x_{n+1} = \sqrt{\cos x_n}$ and the starting value $x_0 = 0.8$ to find the positive root correct to 2 decimal places.

5
$$f(x) \equiv e^{5-2x} - x^5.$$

Show that the equation $f(x) = 0$

- a has a root in the interval (1.4, 1.5),
 b can be written as $x = e^{1-kx}$, stating the value of k .
 c Using the iteration formula $x_{n+1} = e^{1-kx_n}$, with $x_0 = 1.5$ and the value of k found in part b, find x_1, x_2 and x_3 . Give the value of x_3 correct to 3 decimal places.

6

$$f: x \rightarrow 2^x + x^3 - 5, \quad x \in \mathbb{R}.$$

- Show that there is a solution of the equation $f(x) = 0$ in the interval $1.3 < x < 1.4$.
- Using the iterative formula $x_{n+1} = \sqrt[3]{5 - 2^{x_n}}$, with $x_0 = 1.4$, find x_1, x_2, x_3 and x_4 .
- Hence write down an approximation for this solution of the equation $f(x) = 0$ to an appropriate degree of accuracy.

Another attempt is made to find the solution using the iterative formula $x_{n+1} = \frac{\ln(5 - x_n^3)}{\ln 2}$.


- Describe the outcome of this attempt.

7

$$f(x) = 2x^3 + 4x - 9.$$

- Find $f'(x)$.
- Hence show that the equation $f(x) = 0$ has exactly one real root.
- Show that this root lies in the interval $(1.2, 1.3)$.
- Use the iterative formula $x_{n+1} = \sqrt[3]{4.5 - 2x_n}$, with $x_0 = 1.2$, to find the root of $f(x) = 0$ correct to 2 decimal places.
- Justify the accuracy of your answer.

8



The diagram shows part of the curve with equation $y = 3x + \ln x - x^2$ and the line $y = x$. Given that the curve and line intersect at the points A and B , show that

- the x -coordinates of A and B are the solutions of the equation $x = e^{x^2 - 2x}$,
 - the x -coordinate of A lies in the interval $(0.4, 0.5)$,
 - the x -coordinate of B lies in the interval $(2.3, 2.4)$.
- Use the iteration formula $x_{n+1} = e^{x_n^2 - 2x_n}$, with $x_0 = 0.5$, to find the x -coordinate of A correct to 2 decimal places.
 - Justify the accuracy of your answer to part **d**.
- 9
- On the same set of axes, sketch the graphs of $y = x^4$ and $y = 5x + 2$.
 - Show that the equation $x^4 - 5x - 2 = 0$ has exactly one positive and one negative real root.
 - Use the iteration formula $x_{n+1} = \sqrt[4]{5x_n + 2}$, with $x_0 = 1.8$, to find x_1, x_2, x_3 and x_4 , giving the value of x_4 correct to 3 decimal places.
 - Show that the equation $x^4 - 5x - 2 = 0$ can be written in the form $x = \frac{a}{x^3 + b}$, stating the values of a and b .
- Use the iteration formula $x_{n+1} = \frac{a}{x_n^3 + b}$, with $x_0 = -0.4$ and your values of a and b , to find the negative real root of the equation correct to 4 decimal places.

Жартылай бөлу әдісін қолданып табыңыз?

- 1** For each equation, show that it can be rearranged into the given iterative form. Use this and the given value of x_0 to find x_1 , x_2 and x_3 . Give your value of x_3 correct to 4 decimal places.
- a** $9 + 4x - 2x^3 = 0$ $x_{n+1} = \sqrt[3]{2x_n + 4.5}$ $x_0 = 2$
- b** $e^x - 8x + 5 = 0$ $x_{n+1} = \ln(8x_n - 5)$ $x_0 = 3$
- c** $\tan x - 5x + 13 = 0$ $x_{n+1} = \arctan(5x_n - 13)$ $x_0 = -1.2$
- d** $\ln x + \sqrt{x} + 1.4 = 0$ $x_{n+1} = e^{-(\sqrt{x_n} + 1.4)}$ $x_0 = 0.16$
- 2** For each equation, show that it can be rearranged into the given iterative form and state the values of the constants a and b . Use this and the given value of x_0 to find x_1 , x_2 and x_3 . Give your value of x_3 correct to 3 decimal places.
- a** $e^{2x-1} - 6x = 0$ $x_{n+1} = a(\ln bx_n + 1)$ $x_0 = 1.7$
- b** $\frac{2}{x} + \cos x - 3 = 0$ $x_{n+1} = \frac{a}{b - \cos x_n}$ $x_0 = 0.8$
- c** $2x^3 - 6x - 11 = 0$ $x_{n+1} = \sqrt{a + \frac{b}{x_n}}$ $x_0 = 2$
- d** $15 \ln(x + 3) - 4x = 0$ $x_{n+1} = e^{ax_n} + b$ $x_0 = -2.5$
- 3** In each case, use the given iteration formula and value of x_0 to find a root of the equation $f(x) = 0$ to the stated degree of accuracy. Justify the accuracy of your answers.
- a** $f(x) = 10^x + 3x - 4$ $x_{n+1} = \log_{10}(4 - 3x_n)$ $x_0 = 0.44$ 3 decimal places
- b** $f(x) = x^2 + \frac{1}{x-5}$ $x_{n+1} = \sqrt{\frac{x_n^3 + 1}{5}}$ $x_0 = 0.5$ 2 significant figures
- c** $f(x) = 30 - 5x + \sin 2x$ $x_{n+1} = 6 + 0.2 \sin 2x_n$ $x_0 = 6$ 3 significant figures
- d** $f(x) = e^{4-x} - \ln x$ $x_{n+1} = 4 - \ln(\ln x_n)$ $x_0 = 3.7$ 3 decimal places